Rhodora

JOURNAL OF THE

NEW ENGLAND BOTANICAL CLUB

Conducted and published for the Club, by

ALBION REED HODGDON, Editor-in-Chief

ALBERT FREDERICK HILL
STUART KIMBALL HARRIS
RALPH CARLETON BEAN
ROBERT CRICHTON FOSTER
ROLLA MILTON TRYON
RADCLIFFE BARNES PIKE
LORIN IVES NEVLING, JR.
} Associate Editors

VOLUME 68

1966

The New England Botanical Club, Inc.

Botanical Museum, Oxford St., Cambridge 38, Mass.

INDEX TO VOLUME 68

New scientific names and combinations are printed in bold face type

Abies and Cedrus, A New Method for Maintaining the Cones of Intact for Study and Storage 516; balsamea 523; concolor 517 Pl., 523; magnifica 517 Pl.

Acer pensylvanicum 186; spicatum 186

Achillea borealis 414 fig., 417

Aciachne pulvinata 266

Acorus brachystachys 28

Aegopodium podagraria 519

Aegopogon bryophilus 280; cenchroides 280; Fiebrigii 280; geminiflorus, var. muticus 280

Agaricus coloreus 176; lilacinus 181; luteoolivaceus 176; rubescentifolius 176

Agenium villosum 353

Agropyron attenuatum 236; breviaristatum 236; dasystachyum, var. **psammophilum** 437; psammophilum 437

Agrostis alba 438; araucana 259, 261; boliviana 258, 260; exasperata 258, 261; gelida 258, 259; gigantea 438; Haenkeana 259, 260; hyemalis 439; var. tenuis, f. **setigera** 438, 439; michauxii, var. laxiflora 438; montevidensis 258, 259; palustris 438; perennans 258, 260; polymorpha. var. palustris 438; scabra 438, 439; var. scabra, f. tuckermanii 438; var. septentrionalis, f. setigera 438, 439; stolonifera 258, 259; var. compacta 438; var. major 438; var. palustris 437; tolucensis 259, 261; verticillata 259; virginica 448

Aira caryophyllea 515; conferta 243

Alnus tenuifolia 186

Alopecurus aequalis 262; bracteatus 262; Hitchcockii 262

Amaranthus tamariscinus 519

Amphibromus scabrivalvis 115

Andresen, John W. A Multivariate Analysis of the Pinus chiapensis-monticola-strobus Phylad 1

Andropogon altus 348, 350; bicornis 347, 350; cirratus 348, 349; condensatus 348; cordatus 348, 351; gerardii 439; Hassleri 347, 349; hirtiflorus 348, 349; lateralis 348, 351; leptocladus 348, 351; leucostachyus 347, 350; Riedelii 349; saccharoides 348, 350; var. barbinodis 350; ssp. laguroides 350; ssp. parvispiculus 350; subvar. perforatus 350; Selloanus 347, 351; tener 348, 349; subvar. hirtiglumis 348, 349; villosus 353; virgatus 347, 348

Anthochloa lepidula 234; rupestris 234

Aphanelytrum procumbens 115

Aquatic Vascular Plants New for Illinois 221

Arabis canescens 513; Drummondii 413 fig., 417; fendleri, var. spatifolia 513; hirsuta, var. pycnocarpa 413 fig., 417; holboellii, var. retrofracta 513; pauciflora 513

Arctic-Alpine and Boreal Disjuncts, Distribution of; Contributions of the Flora of Nova Scotia VIII 409

Aristida adscensionis 277; Asplundii 278; capillacea 277; circinalis 277, 279; complanata

279; enodis 277, 278; Friesii 279; intermedia 440; inversa 278; implexa 277, 278; laevis 277, 279; longiramea. var. boliviana 279; longispica, var. geniculata 440; Mandoniana 277; mendocina 277, 278; necopina 440; Pflanzii 278; riparia 277, 278
Arnica chionopappa 412 fig., 417, 418
Arundinaria Herzogiana 109
Arundinella Berteroniana 291; confinis 291
Arundo Donax 231
Asplenium callipteris 466; limbatum 467; marginatum 467; Trichomanes 413 fig., 415, 417; viride 410, 411 fig., 416
Asterella **evansii** 470, 471 fig., 472 fig., 473 fig.; from Mexico, A New Species of 470; macropoda 473; venosa 472, 473
Athrostylidium racemiflorum 108
Athyrium filix-femina 127
Avena barbata 244; fatua 244; mollis 449
Axonopus affinis 300; andinus 299, 303; aureus 305; barbigerus 299, 302; capillaris 299, 304; chrysites 299, 305; chrysoblepharis 299, 304; compressus 299, 301; var. macropodius 299, 301; elegantulus 299, 303; eminens, var. bolivianus 299, 302; exasperatus 299, 304; fissifolius 299, 301; Herzogii 299, 304; hirsutus 299, 300; iridifolius 299, 303; marginatus 299, 300; pilosus 298, 302; Purpusii 299, 300; scoparius 299, 303; siccus 299, 302

Ballota alba 94; foetida 93; nigra 93, 94; in the Eastern United States, Varieties of 93; ssp. foetida 93, 94; var. alba 94; var. foetida 93; ssp. nigra 93; var. ruderalis 93
Banks, D. J., Paspalum minus (Gramineae) in Louisiana and Mississippi 94
Barr, Margaret E. and H. E. Bigelow, Contribution to the Fungus Flora of Northeastern North America, IV 175
Beaman, J. H. and C. E. Seidenschnur, Cuchumatanea, A New Genus of the Compositae (Heliantheae) 139
Betula borealis 412 fig., 415, 416; glandulosa 412 fig., 415, 416; Michauxii 414 fig., 416, 417; pumila 414 fig., 417
Berberis Thunbergii 523; vulgaris 523
Bigelow, Howard E. and Margaret E. Barr, Contribution to the Fungus Flora of Northeastern North America, IV 175
Bolivia, Studies in the Flora of, IV., Gramineae 97, 223, 241
Boreal Disjuncts, Distribution of Arctic-alpine and, Contributions to the Flora of Nova Scotia 409
Bouteloua aristidoides 287; curtipendula 287; megapotamica 287; simplex 287
Brachiaria plantaginea 298
Brachypodium mexicanum 112
Britton, D. M. and Rolla Tryon, A Study of Variation in the Cytotypes of Dryopteris spinulosa 59
Briza Mandoniana 223; var. Herzogiana 223; var. tuberculata 223; var. vallegrandensis 223; paleapilifera 223; spicigera 224; stricta 223, 224; uniolae 224

Bromus angustatus 111; arenarius 168, 169; arvensis 170, 173; brizaeformis 170, 173; Buchtienii 111; catharticus 110, 111; commutatus 168, 169, 171, 173, 515; f. commutatus 171, 173, 515; f. **perglaber** 515; f. **pubens** 171, 173; var. pubens 168, 171, 173; lanatus 111; latiglumis 440; lepidus 169, 170, 172; f. lasiolepis 170, 172; f. lepidus 170, 172; f. micromollis 172; Mandonianus 243; mollis 168, 169, 171; and Allies in New England 168; f. leiostachya 169, 171; f. mollis 169, 171; mutabilis, var. hirtus 172; Pflanzii 111; pitensis 111; pratensis 173; pubescens, f. **glaberiflorus** 440; purgans, f. glabriflorus 440; racemosus 168, 170, 173; segetum 110, 111; secalinus 169, 170, 172, 515; var. hirsutus 170, 172, 173; var. secalinus 170, 172; f. **hirtus** 170, 172; f. secalinus 170, 172; f. submuticus 170, 172; Thominii 168, 170, 171; f. hirsutus 170, 171; f. Thominii 170, 171; Trinii 110, 111; unioloides 111

Buchloë dactyloides 36

Burch, Derek, The Application of the Linnaean Names of Some New World Species of Euphorbia subgenus Chamaesyce 155

Burk, John C. and Chung-Hie Nah, Environmental Control of a Taxonomic Character Separating Heterotheca s. str. from Chrysopsis 406

Calamagrostis amoena 246, 249; Antoniana 247, 254; Beyrichiana 256; boliviensis 249; brevifolia 247, 253; calderillensis 246, 248; cephalantha 247, 252; chrysantha 246, 250; cryptolopha 246, 251; curta 247, 252; curvula 246, 252; densiflora 247, 257; elegans 248; eminens 246, 248; var. sordida 248; var. tunariensis 248; Fiebrigii 247, 254; filifolia 249; glacialis 247, 251; heterophylla 247, 249; Humboldtiana 255; intermedia 246, 255; Jamesonii 247, 251; leiophylla 247, 248; Lilloi 257; longiaristata 246, 256; minima 247, 255; montevidensis 257; var. linearis 256; Orbignyana 247, 255; ovata 246, 250; Pflanzii 250; pentapogonoides 250; polygama 246, 257; recta 247, 256; rigescens 246, 249; rosea, var. macrochaeta 253; spicigera 246, 258; tarijensis 246, 254; tarmensis 247, 253; toluccensis, var. laxiflora 248; trichophylla 247, 256; variegata 252, 444; vicunarum 247, 250; violacea 247, 253; viridiflavescens 246, 257

Calamovilfa longifolia, var. magna 437

Callistosporium luteoolivaceum 176, 178; psilocybe 176

Calopogon pulchellus 460; tuberosus 460

Candolleodendron brachystachum 429, 430 fig.; A New Genus of the Leguminosae (Cesalpinioideae) 429

Capparidaceae, Studies in the, VIII Polanisia dodecandra (L.) DC. 41

Carex abdita 453; acuta 451; atratiformis 413 fig., 417; bebbii 451; capillaris 412 fig., 415, 416; var. major 412 fig., 416; communis, f. **gynandra**

451; var. gynandra 451; houghtoniana 451; houghtonii 451, 452; hystericina 452; hystricina 452; lagopodioides 451; lanuginosa 452; lasiocarpa, var. latifolia 452; leptalea, ssp. harperi 452; var. harperi 452; livida, var. grayana 452; var. radicaulis 452; pensylvanica, f. androgyna 451; rugosperma, var. **tonsa** 453; scirpoidea 411 fig., 416; scoparia 451; tribuloides, var. bebbii 451; umbellata, var. tonsa 453

Cedrus and Abies, A New Method for Maintaining the Cones of Intact for Study and Storage 516; atlantica 517 Pl.

Cenchrus echinatus 342; myosuroides 342; pauciflorus 342, 343; viridis 342

Chaetochloa lutescens 446

Chaetosphaeria phaeostroma, var. phaeostromoides 190; phaeostromoides 190

Chamaesyce jenningsii 160; nirurioides 160

Chascolytrium spicigerum 224

Chenopodium graveolens 523

Chloris Beyrichiana 284, 285; distichophylla 284, 285; halophila 285; polydactyla 285; radiata 285, 286; virgata 285; verticillata 518

Chromosome numbers in Vernonia (Compositae) 49

Chromosome numbers in Zinnia and Sanvitalia, New 215

Chrysopsis berlandieri 406; Environmental Control of a Taxonomic Character Separating Heterotheca s. str. from 406

Chusquea delicatula 109, 110; longipendula 110; peruviana 109, 110; quitensis, var. patentissima 110; scandens 110; serrulata 109, 110; spicata 110; uniflora 109, 110

Cirsium altissimum 3; discolor 3

Cladrastis lutea 523

Cleome aspera 44; dodecandra 41, 42, 43, 44, 45, 46, 47; gynandra 43; icosandra 43; monophylla 43; polygama 46; serrata 46; viscosa 43, 44, 46

Clewell, Andre F., Native North American Species of Lespedeza (Leguminosae) 359

Clitocybe **albimontana** 178, 179; asterospora 179

Coix lachryma-jobi 358

Collomia linearis 523

Collybia colorea 176; var. rubescentifolia 176; luteoolivacea 176; rubescentifolia 176

Convallaria canaliculata 458

Copromanthus herbaceus, β ecirratus 459

Corallorhiza innata 460; integra 461; ruppii 460, 461; trifida 460, 461

Cornucopiae hyemalis 438

Correll, Donovan S., Some Additions and Corrections to the Flora of Texas III 420

Cortaderia bifida 231, 232; var. grandiflora 232; rudiuscula 231, 232

Cowan, Richard S., Candolleodendron, A New Genus of Leguminosae (Cesalpinioideae) 429

Cryptogramma Stelleri 413 fig., 417

Cuchumatanea A New Genus of Compositae (Heliantheae) 139; **steyermarkii** 139, 140 fig., 143, 144

Cuphea carthagenensis 423; petiolata 423, 424

Cycloloma atriplicifolium 519

Cynodon Dactylon 283

Cypripedium × Andrewsii 53,

56, 57, 461, 462; Calceolus, var. parviflorum 53, 54, 55, 56, 57; candidum 53, 54, 55, 56, 57, 461; × favillianum 462; × landonii 462; in Manitoba, Natural Hybrids of Lady's Slippers 53
Cystopteris fragilis, var. laurentiana 413 fig., 417

Dactyloctenium aegyptium 283
Danthonia cirrhata 245
Del Rosario, R. M., A New Species of Asterella from Mexico 470
Deschampsia caespitosa 243, 244, 440; conferta 243; elongata 523
Desmodium ciliare 363
Deyeuxia anthoxanthum 250; capitata 250; gracilis 255; imberbis 249; Lagurus 252; leiopoda 250; Mandoniana 255; nivalis 250; obtusata 258; phalaroides 250; picta 251; polystachya 248; subsimilis 258; sulcata 256
Diapensia lapponica 412 fig., 416
Dielsiochloa floribunda 243
Digitaria adscendens 294; horizontalis 294, 295; leiantha 294, 295; lanuginosa 294, 295; violascens 294, 295
Diplazium callipteris 466; celtidifolium 466; **legalloi** 466; **limbatum** 467; marginatum 467
Dissanthelium calycinum 241, 242; macusaniense 241; minimum 241; peruvianum 241; Trollii 241, 242
Distichlis humilis 230; spicata 230; var. andina 230; γ marginata 230
Draba arabisans 413 fig., 417; norvegica 410, 412 fig., 416; pycnosperma 412 fig., 417
Dryopteris × Boottii 90; consimilis 468; cristata, var. Clintoniana × spinulosa, var. intermedia 91; cristata × spinulosa 90; × spinulosa, var. intermedia 90, 91; dilatata 127; fragrans, var. remotiuscula 413 fig., 417; Goldiana × marginalis 90; linnaeana, f. glandulosa 133, 134; marginalis × spinulosa, var. intermedia 90; mollicella 468; spinulosa 59, 60, 67, 69. 70, 76, 77, 78 fig., 79 fig., 81, 82 fig., 84; A Study of Variation in the Cytotypes of 59; var. intermedia 59, 60, 61 fig., 63, 64, 65, 66, 67, 68 fig., 69, 70, 71, 72 fig., 73, 74, 75, 76, 77, 78 fig., 79 fig., 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92; × var. spinulosa 62 fig., 64, 67, 68 fig., 69, 70, 71, 73 fig., 75, 76, 77, 78 fig., 79 fig., 80, 81, 82, 84, 85, 86, 87, 88, 90, 91, 92; var. spinulosa 59, 60, 63 fig., 64, 65, 66, 67, 68 fig., 69, 70, 71, 73, 74 fig., 75, 76, 77, 78 fig., 79 fig., 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92
Duncan, W. H. and S. B. Jones, Jr., Chromosome Numbers in Vernonia (Compositae) 49

Echinochloa crus-galli, var. frumentacea 440, 441; crus-pavonis 335; frumentacea 441; muricata 440
Elaphoglossum dussii 467; petiolatum, var. **dussii** 467
Elocharis calva 453; erythropoda 453; palustris 453
Eleusine indica 282, 515
Elymus angulatus 236
Elyonurus adustus 355, 356;

tripsacoides 355, 356; var. brevidentatus 356; var. ciliaris 356

Emex spinosa 421

Epilobium Hornemanni 414 fig., 417

Eragrostis articulata 225, 227; var. glabrescens 227; bahiensis 224, 228; boliviensis 228; Buchtienii 228; cilianensis 224, 226, 441; ciliaris 224, 225; expansa 225, 229; glomerata 225, 226; hypnoides 225; intermedia 518; lugens 225, 229; lurida 225, 228; maypurensis 225, 226; mexicana 225, 228; montufari 225, 228; nigricans 225, 227; pectinacea 515; pilosa 225, 227, 515; polytricha 225, 229; soratensis 229; spectabilis, var. sparsihirsuta 516; subatra 227; uniolae 224

Erianthus angustifolius 347; Trinii 347

Erigeron hyssopifolius 414 fig., 417

Eriochloa contracta 518; distachya 297; gracilis 518; punctata 297, 298

Eriochrysis cayennensis 346; Warmingiana 346

Erythronium americanum 420

Euphorbia brasiliensis 160; chamaesyce 155, 163, 164; glomerifera 160; hypericifolia 155, 159, 160, 161, 162, 163; hyssopifolia 160, 161, 163; lasiocarpa 160, 161, 163; maculata 155, 156, 157, 158, 159, 160, 161, 164, 165; nutans 160, 163; pilulifera 160; preslii 160, 163; prostrata 155, 158, 163, 164; subgenus Chamaesyce, The Application of Linnaean Names of Some New World Species of 155; thymifolia 155, 158, 164

Ferns, Gymnocarpium, New Data on North American Oak 121; Notes on Lesser Antillean II 464

Festuca arundinacea 441; Buchtienii 114; dolichophylla 113, 114; elatior 441; Fiebrigii 113; laeteviridis 114; megalura 112, 113; orthophylla 113, 114; var. boliviana 114; Pflanzii 114; pratensis 441; procera 112, 113; prolifera 413 fig., 417; var. lasiolepis 412 fig., 417; rigescens 113, 114; sublimis 112, 113; ulochaeta 112, 113

Fore, Paul L. and R. H. Mohlenbrock, Two New Naiads From Illinois and Distributional Records of the Naiadaceae 216

Foster, R. C., Studies in the Flora of Bolivia, IV. Gramineae 97, 223, 241

Frankenia **Johnstonii** 424

Fungus Flora of Northeastern North America, IV., Contribution to the 175

Gentiana × billingtonii 213; × curtisii 213; gracilis 212; linearis 211; puberula 209, 210 Pl., 211, 212, 213, 214; **puberulenta** 210 Pl., 211, 214; A Known But Unnamed Species of the North American Prairies 209; × Gentiana alba 213; × Gentiana andrewsii 213; rigida 212; rubricaulis 211, 212; saponaria 211, 212; *β* linearis 211, 212; shortiana 212; torreyana 212

Ginkgo biloba 521, 523

Glyceria fernaldii 445; pallida, var. fernaldii 445

Gouinia brasiliensis 281; latifolia

281, 282
Gramineae, Studies in the Flora of Bolivia, IV 97, 223, 241
Grammitis limbata 468; marginella 468; **phlegmaria** 467; var. **antillensis** 467; **stipitata** 467
Grear, John W., Jr., Cytogeography of Orontium aquaticum (Araceae) 25
Gross, A. T. H., G. A. Stevenson and H. H. Marshall, Natural Hybrids of Lady's Slippers (Cypripedium) in Manitoba 53
Gymnocarpium continentale 128 129, 135; dryopteris 121, 124 fig., 125, 126, 127 fig., 128, 131, 132, 133, 134, 135, 136, 137; var. glandulosa 134; var. disjunctum 122 Pl., 128, 129, 130 Pl., 135; × var. dryopteris 131; dryopteris, var. pumilum 134; var. dryopteris 128, 129, 136; × robertianum 133; **heterosporum** 123 Pl., 124 fig., 127 fig., 130 fig., 131, 132, 134, 135, 136, 137; New Data on North American Oak Ferns 121; robertianum 121, 125, 126, 127 fig., 128, 131, 133, 134, 135, 136
Gymnocladus dioica 523
Gymnogramma gracilis 468
Gymnopogon jubiflorus 284; spicatus 284
Gynerium sagittatum 232

Hackelochloa granularis 357
Hedysarum conglomeratus 371; divergens 390; frutescens 370, 381, 390, 392; hirtum 376, 377; prostratum 385; repens 385, 386; reticulatum 393; sessiliflorum 393; umbellatum 370; violaceum 380, 390, 392
Hehre, Edward J., Polanisia dodecandra in New Hampshire 48
Helianthemum tripetalum 208
Hemerocallis flava 456; lilio-asphodelus 456, 457
Hemidictyum marginatum 467
Herbarium, Botanical Notes from the Pringle, I 514
Heteropogon contortus 353, 354; melanocarpus 353, 354; villosus 353
Heterotheca latifolia 406; var. macgregoris 406; s. str., Environmental Control of a Taxonomic Character Separating Chrysopsis from 406; subaxillaris 406, 407, 519; var. petiolaris 406
Hodgdon, A. R., Revised Checklist of the Vascular Plants of Maine (Review) 521; and Frederic Steele, Rubus Subgenus Eubatus in New England 474
Holcus lanatus 244; sorghum 441
Homolepis aturensis 334
Hordeum halophilum 237; muticum 237; nodosum 237
Hounsell, R. W. and E. C. Smith, Contributions to the Flora of Nova Scotia VIII. Distribution of Arctic-alpine and Boreal Disjuncts 409
Hyacinthus muscari 457, 458; racemosus 457, 458
Hygrophorus lilacinus 179, 181; monticola 180 Pl., 181, 182; nitiosus 182; tahquamenonensis 182; violeïpes 181
Hymenachne donacifolia 333
Hymenophyllum gratum 464, 465; hirsutum 465, var. **gratum** 465; hirtellum 464, 465; var. gratum 465, 466; var. vincentinum 465; maxonii 465
Hyparrhenia bracteata 351

Ichnanthes breviscrobs 329, 330; calvescens 329, 330; candicans 330, 331; minarum 330, 331; pallens 330, 331; peruviana 329, 330; Ruprechtii 330, 331
Illinois, Aquatic Vascular Plants New for 221
Iltis, Hugh H., Studies in the Capparidaceae VIII. Polanisia dodecandra (L.) DC. 41
Imperata brasiliensis 345; minutiflora 345; tenuis 345
International Plant Index, Authors of Plant Genera and the (Review) 35
Isachne arundinacea 334

Jaegeria discoidea 145; gracilis 145; hirta 140 fig., 143, 144, 145
Jones, S. B., Jr. and W. H. Duncan, Chromosome Numbers in Vernonia (Compositae) 49
Juglans nigra 523
Juncus alpinus, var. rariflorus 414 fig., 417; campestris β 455; congesta 454; erectus 454; glabratus 454; intermedius 454, 455, 456; multiflorus 454, 455, 456; trifidus 411 fig., 416; validus 519
Juniperus communis, var. communis 514
Jussiaea diffusa 218

Keissleriella caudata 185, 187 fig., cladophila 185
Koeleria cristata 242, 441; gracilis, ssp boliviensis 242; macrantha 441; pseudocristata, var. andicola 242

Lactuca muralis 138; in New England 138
Lamprothyrsus Hieronymi 245; var. tinctus 245
Lasiacis ligulata 332; sorghoidea 332
Lasiosphaeria chrysentera 185; ovina 185
Lechea brevifolia 205; cernua 203; cinerea 194; (Cistaceae), Notes on Rafinesque's Species of 192; corymbosa 200; divaricata 203; deckertii 203; floridana 203, 204; furfuracea 195; glomerata 200; heterophylla 199, 200; var. major 200; var. minor 200; var. parviflora 200; intermedia 193, 195, 198; juncifolia 196, 197; laxiflora 195, 196, 197; var. brevifolia 196; leggettii 193, 194, 195, 197, 198, 201, 203, 207; linifolia 208; major 197; maritima 193, 198, 205; mexicana 208; minor 194, 198, 199, 200, 201, 202, 203, 204, 205, 206, 207; mucronata 197, 198, 200, 201, 202, 203, 204; var. ramosa 199; var. sessiliflora 199; var. simplex 199; var. typica 199; paucifolia 208; pauciflora 205; patula 194, 203, 204, 207; pulchella 193, 194, 196, 197, 203; racemulosa 193, 195, 197, 198, 201, 202, 203, 204; recurvata 197; revoluta 202; secundiflora 204; sessiliflora 207, 208; stellata 206, 207; stricta 195; surculosa 201, 202; tenuifolia 196, 197, 198, 199, 200, 202, 203, 204, 205; ternifolia 194, 195; torreyi 194, 203, 207; thymifolia 194, 205, 206; uniflora 206; verna 196, 197; villosa 197, 198, 199, 200, 202, 203, 204, 207; var. macrotheca 199; var. schaffneri 199; virgata 202, 203; var. bracteata 203; var. breviflora 203
Ledum groenlandicum 189

Leiosphaerella **falcata** 186, 187 fig., phoenicis 186; praeclara 186

Leptochloa domingensis 280, 281; dubia 280, 281; uninervia 280, 519; virgata 280, 281

Leptocoryphium lanatum 292

Leptoloma cognatum 516

Lespedeza acuticarpa 403; angustifolia 363, 367, 369, 370, 372, 375 fig., 383, 393, 394, 395, 399; × capitata 396, 400 fig.; × hirta 364, 377, 396, 398, 399, 401 fig.; × repens 396, 400 fig.; × stuevei 396, 400 fig.; × virginica 396; var. brevifolia 368, 369; f. subvelutina 368, 369; bicknellii 371; bicolor 359; brittonii 402; capitata 362, 363, 367, 370, 372, 373, 374, 375 fig., 379, 382, 393, 394, 398, 399; × hirta 396, 398, 399, 401 fig.; × intermedia 396, 397, 400 fig.; × leptostachya 396, 401 fig.; × repens 396, 401 fig.; × stuevei 396, 400 fig.; violacea 396, 399, 400 fig.; virginica 396, 397, 399, 401 fig.; var. angustifolia 368; var. calycina 376, 399; var. capitata 373; var. hirtiformis 398, 399; var. longifolia 399; var. sericea 371, 373; var. stenophylla 371, 373, 374; f. argentea 371, 374; var. typica 371, 373; var. velutina 371, 373; var. vulgaris 371, 374; cuneata 359; divergens 390, 393; frutescens 371, 380, 381, 392, 393; var. acutifructa 380, 381; hirta 363, 367, 368, 372, 374, 377, 378, 379, 399; × intermedia 396, 397, 399, 400 fig., 402; × procumbens 396, 401 fig.; × repens 396, 403; × stuevei 396, 397, 401 fig.; × violacea 377, 396, 400 fig.; × virginica 396, 397, 403; ssp. curtissii 375 fig., 376, 377, 378, 379, 398; ssp hirta 375 fig., 376, 377, 378; var. angustifolia 368; var. appressipilis 377, 398; var. calycina 376, 379; var. dissimulans 398, 399; var. intercursa 368, 370; var. longifolia 398; var. oblongifolia 398; var. sparsiflora 376, 377; var. typica 376; intermedia 361 fig., 363, 368, 375 fig., 380, 381, 388, 392, 393, 397, 402; × procumbens 396, 401 fig.; × repens 396, 400 fig.; × stuevei 396, 403 fig.; × violacea 396, 400 fig.; × virginica 396, 400 fig.; f. hahnii 402; var hahnii 401; (Leguminosae), Native North American Species of 359; leptostachya 367, 370, 375 fig., 382, 383, 395; longifolia 398, 399; manniana 399; neglecta 402, 403; × nuttallii 397; nuttallii 399; var. manniana 399; var. nuttallii 402; oblongifolia 398; polystachya 376; prairea 390, 393; procumbens 363, 364, 367, 383, 384 fig., 385, 390, 392; × repens 396, 400 fig.; × stuevei 364, 396, 403 fig.; × violacea 396, 400 fig.; × virginica 396, 401 fig., 402; var. elliptica 402; prostrata 385; reticulata 380, 393; var. angustifolia 393; var. sessiliflora 380; repens 361 fig., 367, 384 fig., 385, 386, 387, 390, 392, 393; × stuevei 396, 401 fig.; × violacea 396, 401 fig.; × virginica 396, 403 fig.; sessiliflora 380, 393; simulata 397, 399; stuevei 363, 368, 371, 384 fig., 387, 388, 395, 402; × violacea 396, 401 fig.;

× virginica 396, 402, 403 fig.; f. angustifolia 402, 403; var. angustifolia 402, 403; var. intermedia 380; var. neglecta 402; stipulacea 359; striata 359; texana 367, 389, 390, 391 fig., 395; velutina 371, 375; virgata 401; virginica 363, 368, 381, 388, 391 fig., 393, 394, 395; f. deamii 402, 403; var. deamii 395; var. sessiliflora 380; var. typica 394; violacea 361 fig., 362, 363, 368, 381, 387, 390, 391 fig., 392, 393; × virginica 396, 400 fig., 403; var. angustifolia 393, 394; var. divergens 390; var. prairea 390; var. reticulata 393; var. sessiliflora 380

Limonium limbatum, var. **glabrescens** 425; var. limbatum 425

Linnaea borealis, var. americana 187, 188

Linnaean Names of Some New World Species of Euphorbia subgenus Chamaesyce, The Application of 155

Liriodendron tulipifera 523

Lolium multiflorum 238; perenne 238; temulentum 238; var. arvense 238

Long, Robert W., Observations Regarding the Occurrence and Relationships of Ruellia Lorentziana (Acanthaceae) in Southern Florida 432

Louisiana and Mississippi, Paspalum minus in 94

Luziola peruviana 290

Luzula congesta 456; intermedia 454, 456; multiflora 454, 455, 456; parviflora 414 fig., 417; var. melanocarpa 414 fig., 417; spicata 412 fig., 416, 418

Lycopodium reflexum, var. reflexum 464; var. rigidum 464; rigidum 464; Selago 411 fig., 415, 416

Lycurus phleoides 263

Lysichitum americanum 28; camtschatcense 27

MacDonald, R. D. and M. E. MacDonald, A New Method for Maintaining the Cones of Abies and Cedrus Intact for Study and Storage 516

Magnolia acuminata 523

Maine, Revised Check-list of the Vascular Plants of (Review) 521

Manisuris altissima 356; aurita 356, 357; fasciculata 356

Manitoba, Natural Hybrids of Lady's Slippers (Cypripedium) in 53

Marshall, H. H., A. T. H. Gross and G. A. Stevenson, Natural Hybrids of Lady's Slippers (Cypripedium) in Manitoba 53

Medicago virginica 393

Melica adhaerens 233; var. tenuis 233; Mandonii 233; scabra 233; var. glabra 233; triflora 450; violacea 233

Mesosetum cayennense 297

Mexico, A New Species of Asterella from 470

Microchloa indica 283

Mississippi, Paspalum minus in Louisiana and 94

Mohlenbrock, R. H. and Paul L. Fore, Two New Naiads from Illinois and Distributional Records of the Naiadaceae 216

Monocots, Nomenclatural Notes on 435

Morton, C. V., Varieties of Ballota nigra in the Eastern United States 93

Muhlenbergia angustata 263,

265; asperifolia 264, 265; fastigiata 264, 265; Herzogiana 264; Holwayorum 264, 266; ligularis 264, 265; peruviana 263, 264; phragmitoides 266; quitensis 264; rigida 264, 266; sobolifera, var. setigera 518; tenuissima 263, 264
Munroa andina 288; var. breviseta 288; argentina 288; decumbens 288
Muscari ambrosiaceum 458; atlanticum 457, 458; moschatum 457, 458; muscari 458; muscarimi 458; racemosum 457, 458
Mycena atkinsoniana 182
Mycosphaerella crepidophora 188; ebulina 188; **linnaeae** 187, 188, 189; minor 188
Myriophyllum heterophyllum 425; verticillatum 425

Nah, Chung-Hie and John C. Burk, Environmental Control of a Taxonomic Character Separating Heterotheca s. str. from Chrysopsis 406
Naiads from Illinois and Distributional Records of the Naiadaceae, Two New 216
Naias flexilis 217, 218, 219; gracillima 217, 218, 219; guadalupensis 217, 218, 219; minor 217, 218, 219, 220
Najas marina 221; minor 222
Nassella Asplundii 269; corniculata 270; deltoidea 269; flaccidula 269; var. humilior 269; Meyeniana 269, 270; pubiflora 269
Naucoria kauffmanii 183
Neottia gracilis 462; lacera 462
New England, Lactuca muralis in 138; Rubus Subgenus Eubatus in 474
New Hampshire, Polanisia dodecandra in 48
Nova Scotia, Contributions to the Flora of VIII. Distribution of Arctic-alpine and Boreal Disjuncts 409

Olyra Buchtienii 343, 344; ciliatifolia 343, 344; ecaudata 343, 344; heliconia 343, 344; latifolia 343, 344; lateralis 343; micrantha 343, 344
Omphalia luteolilacina 181
Omphalina oreades 178, 179
Ophrys corallorhiza 460
Oplismenus hirtellus 334
Orchis rotundifolia, f. beckettae 462; f. beckettii 462; f. **lineata** 462; var. lineata 462
Orontium aquaticum 25; (Araceae), Cytogeography of 25; fossile 28
Orthoclada laxa 234
Oryzopsis florulenta 276
Oxyria digyna 410, 412 fig., 416
Oxytropis johannensis 414 fig., 417

Panicum agrostidiforme 443; agrostoides 443; anceps 443; aquaticum 318, 320; var. chloroticum 320; ashei 441, 442; boliviense 318, 323; capillare, var. minus 442; cayennense 319, 320; chloroticum 320; commutatum, var. ashei 441, 442; var. minor 442; cordovense 318, 327; cyanescens 319, 325; depauperatum, var. involutum 443; var. psilophyllum 443; echinulatum 318, 320; var. boliviense 320; frondescens 318, 324; frumentaceum 441, 446; giganteum 322; glutinosum 319, 325; hebotes 319, 327; hirticaule 319, 320; involutum 443; laxum 318, 323, 324;

lutescens 445, 446; maximum 319, 322; megiston 318, 328; muricatum 440; nervosum 319, 325; olyroides 319, 329; pantrichum 318, 327; parvifolium 319, 326; paucispicatum 318, 328; peladoënse 319, 321; pilosum 318, 323; polygonatum 318, 322; procurrens 318, 329; pulchellum 318, 324; quadriglume 317, 319, 321; rigidulum 443; Rudgei 319, 326; sciurotis 319, 327; Sellowii 319, 326; sphaerocarpon, var. inflatum 519; stoloniferum 318, 324; stramineum 319, 321; trichanthum 319, 325; tricholaenoides 318, 322; versicolor 320, 328; virescens 446

Pappophorum mucronulatum 235; Wrightii 235

Pariana bicolor 239, 240; gracilis 239; lunata 240; **Swallenii** 239; zingiberina 240

Parnassia parviflora 414 fig., 417

Paronychia Jamesii, var. Jamesii 423; var. **praelongifolia** 423

Paspalum boliviense 306, 317; Buchtienii 307, 310; candidum 305, 308; Ceresia 306, 309; ciliatifolium 444; var. muhlenbergii 444; collinum 308, 315; conjugatum 307, 313; conspersum 308, 315; decumbens 306, 311; densum 306, 316; depauperatum 306, 309; dilatatum 307, 313; distichum 307, 311; Ekmanianum 307, 316; erianthum 307, 310; Humboldtianum 307, 310; inaequivalve 306, 313; inconstans 307, 311; Juergensii 307, 312; lepidum 307, 312; lineispatha 306, 309; malacophyllum 306, 316; melanospermum 307, 314; membranaceum 309; millegrana 306, 315; minus 94, 95, 96, 306, 314; (Gramineae) in Louisiana and Mississippi 94; muhlenbergii 443; multicaule 307, 312; notatum 95, 306, 314; paniculatum 307, 312; penicillatum 306, 308; pictum 306, 312; pilosum 306, 316; plicatulum 308, 314; polyphyllum 307, 310; prostratum 306, 308; pubescens 443, 444; pygmaeum 306, 308; var. glabrescens 308; remotum 307, 316; saccharoides 306, 317; stellatum 306, 309; tripinnatum 303; Urvillei 307, 314; virgatum 306, 315

Paxillus curtisii 183

Pennisetum chilense 340; latifolium 339, 341; mutilatum 340, 341; sagittatum 341; setosum 339, 340; tristachyum 340, 341; ssp. boliviense 341; villosum 339, 340

Phaeocollybia kauffmanii 183, 184

Phalaris angusta 289; arundinacea 444; f. picta 444, 445; f. variegata 444, 445; canariensis 289

Pharus glaber 290; latifolius 290; parvifolius 290, 291

Phleum alpinum 413 fig., 417; pratense, var. nodosum 516

Phragmites communis 231

Phyllodoce caerulea 412 fig., 416

Picea glauca 523; rubens 523

Pinguicula vulgaris 412 fig., 416 418

Pinus ayacahuite 16; chiapensis 1, 2, 4, 5, 8 Pl., 10 Pl., 11, 14, 15, 16, 20, 21, 23; -monticola-strobus Phylad, A Multivariate Analysis of the 1; contorta 3; flexilis 13; griffithii 16; monticola 4, 8 Pl., 10 Pl., 12, 14, 15, 16, 20, 21, 23; × strobus 23;

oocarpa 14; parviflora 16; peuce 16; strobiformis 13; strobus 1, 2, 4, 5, 7, 8 Pl., 10 Pl., 11, 12, 14, 15, 16, 20, 21, 23; var. chiapensis 1, 2, 4, 5, 14
Piptochaetium panicoides 270; f. subpapillosum 270; setifolium 270; tuberculatum 270
Plagiophiale petrakii 189
Plagiostigma petrakii 189
Plant Genera and the International Plant Index (review) 35
Plantago insularis, var. fastigiata 428; minima 428; scariosa 428
Plectosphaeria salicis 189
Pleurotus elongatipes 184, 185
Poa alpina 411 fig., 416, 418; androgyna 119; annua 116, 119; asperiflora 116, 118; boliviensis 118; Buchtienii 116; var. subacuminata 116; candamoana 116, 120; denticulata 116, 119; dumetorum 119; var. unduavensis 119, 120; glaucantha 411 fig., 416; gymnantha 116, 117; var. aperta 117; Hieronymi 116, 118; horridula 116, 119; humillima 116, 117; Lilloi 116, 118; myriantha 118; palustris 518; perligulata 117; Pflanzii 118; pratensis 116, 118; scaberula 116, 117
Podosphaera clandestina, var. clandestina 189; oxyacanthae, var. oxyacanthae 190
Polanisia dodecandra 41, 42, 46, 47, 48; in New Hampshire 48; (L.) DC., Studies in the Capparidaceae VIII 41; ssp. **trachysperma** 47; ssp. dodecandra 47; graveolens 41, 42, 45, 46; trachysperma 47; uniglandulosa 47
Polygonatum canaliculatum 458; commutatum 458
Polypodium flabelliforme 467; phlegmaria 467; rigescens 467
Polygonum viviparum 189
Polypogon elongatus 262; interruptus 262, 263
Polystichum Lonchitis 413 fig., 417
Poole, James P., Lactuca muralis in New England 138
Potamogeton americanus 218
Primula mistassinica 414 fig., 417
Pringle, James S., Gentiana puberulenta, sp. nov., a Known but Unnamed Species of the North American Prairies 209
Proboscidea **sabulosa** 426, 427 Pl.
Proctor, George R., Notes on Lesser Antillean Ferns II 464
Pseudechinolaena polystachya 296
Psoralea melitoides 392; psoralioides 392
Puccinellia **fernaldii** 445; oresigena 115; parvula 115

Rafinesque's Species of Lechea (Cistaceae), Notes on 192
Rhoades, Mary H., Seed Germination of Shortia galacifolia T & G Under Controlled Conditions 147
Rhododendron canadense 36
Rhynchelytrum repens 296
Rhynchospora capitellata 453; glomerata, var. capitellata 453; var. minor 453
Rogers, Ken E. and J. K. Underwood, Some Plants New to Tennessee 518
Rollins, Reed C., Authors of Plant Genera and the International Plant Index (Review) 35; The Identity of Sisymbrium pauciflorum (Cruciferae) 513
Rubus abactus 484, 510; abbrevians 482, 510; aculiferus 483, 510; adjacens 510; alius 510;

allegheniensis 475. 484, 486, 487, 494, 503, 504, 507, 508, 510, 511; × pensilvanicus 510, 511; alter 510; alumnus 510; amicalis 510; amnicola 510; Andrewsianus 510; apparatus 510; aptatus 510; arcuans 510; arenicola 478, 481, 486, 488, 493, 510. 511; argutus 483, 487, 499, 505, 510, 511; arundelanus 510; ascendens 510; auroralis 510; avipes 510; Baileyanus 479, 480, 510; barbarus 510; bellobatus 484, 510; Bicknellii 481, 510; biformispinus 510; bifrons 487; Bigelovianus 510; Blakei 510; Blanchardianus 510; Bootianus 484, 510; Brainerdii 510; brevipedalis 510; canadensis 477, 484, 486, 500, 502, 508, 511; coloniatus 510; conanicutensis 510; cubitans 510; cuneifolius 477, 487, 499; curtipes 510; Deaneanus 510; eflagellaris 510; electus 510; elegantulus 478, 481, 482, 483, 486, 487, 497, 500, 502, 503, 510, 511; Enslenii 486, 488, 490, 510, 511; facetus 510; felix 510; Fernaldianus 510; flagellaris 480, 482, 484, 486, 487, 488. 489, 490, 491, 492, 510, 511; frondisentis 510; frondosus 480, 481, 486, 487, 505, 506, 507, 508, 510, 511; fruticosus 475; geophilus 510; glandicaulis 510; gnarus 510; Gravesii 510; Groutianus 497, 510; harmonius 510; heterogeneous 510; heterophyllus 510; hispidoides 510; hispidus 475, 483, 486, 493, 494, 510, 511; × allegheniensis 510, 511; × flagellaris 511; × setosus 510, 511, 512; × vermontanus 512; honorus 510; insons 510; insulus 510; invisus 510; jacens 478, 510; jactus 510; Janssonii 510; Jaysmithii 482, 486, 488, 489, 511; Jeckylanus 510; jugosus 510; junceus 510; junior 510; Kennedyanus 510; laciniatus 487; laevior 510; latens 511; Lawrencii 511; licitus 511; longissimus 511; mainensis 511; maniseesinsis 511; Millspaughii 511; miscix 511; montpelierensis 511; multiformis 511; multilicius 511; multispinus 507, 511; nigricans 511; nigrobaccus 511; notatus 511; novanglicus 511; obsessus 511; orarius 484, 511; ortivus 511; ostryifolius 484, 511; paludivagus 483. 511; Parlinii 511; paulus 511; pauper 511; peculiaris 511; pensilvanicus 481, 484, 487, 501, 503, 505, 507, 508, 510, 511; pergratus 481, 511; perinvisus 511; permixtus 511; perpauper 511; pervarius 511; philadelphicus 511; plicatifolius 479, 481, 492, 511; positivus 511; procumbens 511; prosper 511; pudens 511; pugnax 511; Randii 511; ravus 511; recurvans 481, 511; recurvicaulis 478, 479, 481, 484, 486, 488, 489, 491, 492, 493, 510, 511, 512; × setosus 510; × vermontanus 511; rhodinsulanus 511; rixosus 511; roribaccus 511; Rosa 511; Rossbergianus 511; saltuensis 511; Sanfordii 511; sativus 511; scambens 511; sceleratus 480, 511; semierectus 511; semisetosus 478, 483, 486, 497, 498, 499, 510, 511; sempervirens 511; setosus 475, 483, 484, 486, 494, 495, 497, 499, 502, 504, 510, 511, 512; × al-

legheniensis 510, 511; × vermontanus 510, 511, 512; severus 480, 497, 511; significans 511; singulans 511; spiculosus 511; Subgenus Eubatus in New England 474; tardatus 478. 511; tetricus 511; tholiformis 511; trifrons 512; udus 512; usus 512; univocus 512; vermontanus 478, 481, 484, 486, 494, 496, 497, 498, 499, 500, 502, 510, 511; × allegheniensis 511; × elegantulus 511; × pensilvanicus 511; vigoratus 512; viridifrons 512
Ruellia caroliniensis 432; intermedia 434; lorentziana 433 Pl.. 434; (Acanthaceae) in Southern Florida, Observations Regarding the Occurrence and Relationships of 432; nudiflora 434; var. occidentalis 432; occidentalis 432; succulenta 432; tuberosa 434
Ruppia maritima 221

Saccharum officinarum 345
Sacciolepis myuros 333
Sanvitalia abertii 215; fruticosa 216; New Chromosome Numbers in Zinnia and 215; tenuis 215, 216; versicolor 216
Saxifraga aizoides 410, 412 fig., 416; Aizoon, var. neogaea 409, 412 fig., 416
Schizachne purpurascens 413 fig., 417
Sciadopitys verticillata 521, 523
Scirpus caespitosus, var. delicatulus 413 fig., 417
Scrophularia marilandica 425
Sedum Rosea 411 fig., 415, 416
Seidenschnur, C. E. and J. H. Beaman, Cuchumatanea, A New Genus of the Compositae (Heliantheae) 139
S lloa plantaginea 144
Senecio pauperculus 414 fig.. 417
Sesuvium **erectum** 421, 422; maritimum 422; Portulacastrum 422, 423; sessile 422; **trianthemoides** 422; verrucosum 421
Setaria argentina 336, 338; barbinodis 336, 339; geniculata 336; glauca 445, 446; gracilis 336; f. brevispica 336; f. penicillata 336; Liebmannii, f. trichorachis 339; muricata 440; oblongata 336, 338; Poiretiana 336; scandens 336, 337; tenacissima 336, 337; tenax 336, 337; trichorachis 336, 339; vulpiseta 336, 338
Seymour, Frank C., Botanical Notes from the Pringle Herbarium, I. 514; Bromus mollis and Allies in New England 168
Shortia galacifolia 147, 151, 152, 154; T & G Under Controlled Conditions, Seed Germination in 147
Siegesbeckia repens 144
Sisymbrium pauciflorum 513; (Cruciferae), the Identity of 513
Sisyrinchium angustifolium 459; arenicola 459; atlanticum 459; farwellii 459
Smilax ecirrata 459
Smith, C. E., Jr., Identity of Witchweed in the Southeastern United States 167
Smith, E. C. and R. W. Hounsell, Contributions to the Flora of Nova Scotia VIII. Distribution of Arctic-alpine and Boreal Disjuncts 409
Smith, Gary L., Spergula morisonii in America 520
Solidago multiradiata 412 fig., 416

Sorghastrum minarum 352; parviflorum 352, 353; stipoides 352, 353
Sorghum bicolor 446; halepense 352; vulgare 352, 446
Sparganium chlorocarpum, f. **acaule** 436; simplex, var. acaule 436
Spartina michauxiana, var. suttiei 446; pectinata, var. suttiei 446
Spergula morisonii 520; in America 520; pentandra 520
Sphaerella eucarpa 189; leightonii 188
Sphaeria phaeostromoides 190
Spiranthes gracilis 462, 463; lacera 462, 463
Sporobolus aeneus 267, 268; argutus 267; Berteroanus 268; cryptandrus 518; eximius 269; indicus 267, 268; neglectus 448; Poiretii 267, 268; tenuissimus 267; vaginiflorus 448; var. neglectus 447; virginicus 448
Steele, Frederic, and A. R. Hodgdon, Rubus Subgenus Eubatus in New England 474
Stipa boliviensis 275; brachyphylla 271, 275; capilliseta 271, 273; curviseta 271, 275; dasycarpa 271, 275; depauperata 271, 274; florulenta 271, 276; Hans-meyeri 271, 273; Holwayi 271, 276; Ichu, 271, 272; illimanica 272, 276; inconspicua 271, 274; mucronata 271, 273, 274; nardoides 271, 273; Neesiana 271, 274; obtusa 271, 275; pampagrandensis 271, 272; Pflanzii 269; plumosa 271, 272, 276; polyclada 271, 275, 276; speciosa 271, 272; tiraquensis 269; Trollii 272, 276
Stevenson, G. A., H. H. Marshall, and A. T. H. Gross, Natural Hybrids of Lady's Slippers (Cypripedium) in Manitoba 53
Striga asiatica 167; lutea 167
Stylagrostis leiopoda 250
Swartzia brachystachya 429
Symplocarpus foetidus 27
Syringa vulgaris 523

Targionia hypophylla 471, 472
Tennessee, Some Plants New to 518
Texas, Some Additions and Corrections to the Flora of, III 420
Thaxteria fusca 190
Thelypteris gracilis 468; **consimilis** 468; **cooleyi** 468, 469; limbata 469; noveboracensis 125; palustris 124; piedrensis 468
Thrasya campylostachya 295
Tofieldia glutinosa 414 fig., 417
Torres, Andrew M., New Chromosome Numbers in Zinnia and Sanvitalia 215
Torreyochloa fernaldii 445; pallida, ssp. pallida, var. fernaldii 445
Trachypogon canescens 355; montufari 354, 355; plumosus 355; polymorphus, var. bolivianus 355
Tragus racemosus 279
Trichachne affinis 293; californica 293; insularis 293; saccharata 294; sacchariflora 293
Trichloris mendocina 286; pluriflora 286
Trichodium laxiflorum 438
Tricholaena rosea 296
Tricholoma rubescentifolium 176
Trichopteryx flammida 291
Triglochin maritimum 436; palustre 436
Triniochloa stipoides 266

Triodia avenacea 235
Triphora trianthophora, var. Schaffneri 420; var. trianthophora 420
Triplasis purpurea 518
Tripogon spicatus 282
Tripsacum australe 357; dactyloides 357
Trisetum floribundum 243; oreophilum 243; spicatum 243, 449; var. maidenii 449, 450; var. molle 411 fig., 416, 449; var. pilosiglume 411 fig., 416, 449; ssp molle 450; subspicatum, f. maidenii 450; triflorum. ssp. molle 450
Tristachya chrysothrix 292
Tryon, Rolla and Donald M. Britton, A Study of Variation in the Cytotypes of Dryopteris spinulosa 59
Underwood, J. K. and Ken E. Rogers, Some Plants New to Tennessee 518
Utricularia cornuta 428; juncea 428

Vaccinium angustifolium 189; boreale 414 fig., 417; caespitosum 414 fig., 417; ovalifolium 414 fig., 417; uliginosum 189; var. alpinum 412 fig.. 415, 416
Vernonia acaulis 50; altissima 51; angustifolia 51; var. angustifolia 51; var. mohrii 51; var. scaberrima 51; anthelmintica 51; baldwinii 49, 51; blodgettii 51; cinerea 49, 51; (Compositae), Chromosome Numbers in 49; colorata 51; conferta 51; × concinna 51; ×dissimilis 51; glabra 51; greggii 51; guadaloupensis 51; guineensis 51; interior 51; karvinskiana 51; kotschiana 51; lindheimeri 51; marginata 51; noveboracensis 52; nudiflora 52; ovalifolia 52; perottetii 52; platensis 52; primulina 52; pulchella 52; × recurva 52; texana 52
Vilfa vaginiflora 447, 448
Voss, Edw. G., Nomenclatural Notes on Monocots 435

Wagner, W. H., Jr., New Data on North American Oak Ferns, Gymnocarpium 121
Wilbur, Robert L., Notes on Rafinesque's Species of Lechea (Cistaceae) 192
Winterringer, Glen S., Aquatic Vascular Plants New for Illinois 221
Witchweed in the Southeastern United States, Identity of 167
Woodsia alpina 411 fig., 416; glabella 411 fig., 416; ilvensis 411 fig., 415, 416

Zannichellia palustris 222
Zea mays 167
Zeugites mexicana 234; var. glandulosa 234
Zinnia maritima 215; purpusii 215; and Sanvitalia, New Chromosome Numbers in 215